



FREE NALOXONE

HOW TO RECOGNIZE AN OVERDOSE

The following are signs and symptoms to look for to identify a potential overdose:

LOSS OF CONSCIOUSNESS, LIMP BODY	SLOWED BREATHING, CHOKING	SMALL, "PINPOINT" PUPILS	PALE, BLUE OR COLD SKIN	SLOWED HEART RATE

HOW TO USE NARCAN NASAL SPRAY:

Before use, check for signs of opioid overdose:

- Will not wake up or respond to your voice or touch
- Breathing is very slow, irregular, or has stopped
- Center part of their eye is very small, sometimes called "pinpoint pupils"

Lay the person on their back to receive a dose of NARCAN Nasal Spray.



- 1** Peel back the package to remove the device, hold the device with your thumb on the bottom of the plunger and 2 fingers on the nozzle.
- 2** Tilt the person's head back and provide support under their neck with your hand. Gently insert the tip of the nozzle into one nostril. Until your fingers touch the bottom of the person's nose.
- 3** Press the plunger firmly to give the dose of NARCAN Nasal Spray into the patient's nose. Remove the NARCAN Nasal Spray from the nostril after giving the dose. Move the person on their side (recovery position) after giving NARCAN Nasal Spray.
- 4** Get emergency medical help right away. Watch the person closely. If the person does not respond by waking up to voice or touch, or breathing normally, another dose may be given. NARCAN Nasal Spray may be dosed every 2 to 3 minutes, if available. Repeat Step 2 using a new NARCAN Nasal Spray to give another dose in the other nostril. If additional NARCAN Nasal Sprays are available, repeat step 2 every 2 to 3 minutes until the person responds or emergency medical help is received.

Use NARCAN Nasal Spray (naloxone hydrochloride) for known or suspected opioid overdose in adults and children. Important: For use in the nose only. Do not remove or test the NARCAN Nasal Spray until ready to use.

Window: 18.75 x 17"

Plastic paper (inside): 17 x 12.5"

6"

FREE NALOXONE

8.25"

Sides bottom
Size: 6 x 8.25"

NALOXONE

HOW TO RECOGNIZE AN OVERDOSE

The following are signs and symptoms to look for to identify a potential overdose:

- Loss of consciousness, limp body
- Slowed breathing, choking
- Small, "pinpoint" pupils
- Pale, blue or cold skin
- Slowed heart rate



AEK
Health Supply Company

CONTACT FOR REFILL:
Schuylkill County Mental Health, Developmental Services and Drug & Alcohol Programs
410 N. Centre St. Suite 2, Pottsville, PA 17901
Phone: 570-621-2890
Crisis Line: 1-877-993-4357



11.5"

12.5"

Qty. 2
(4 stickers)

Decals for both side panels
Size: 11.5 x 12.5"

BEWARE OF COUNTERFEIT DRUGS THAT MAY CONTAIN FENTANYL

Learn to Administer Naloxone FAST! TRAINING VIDEOS



AEK
Health Supply Company

Decals for lower front panel
Size: 9.5 x 10"

9.5"

10"